




## Product Specification Sheet


<b>Product Name</b>	Stemgent® TRA-1-60 Antibody (Purified), Mouse anti-Human
<b>Description</b>	The TRA-1-60 monoclonal antibody reacts with a pluripotent-stem-cell-specific antigen expressed on undifferentiated human embryonic stem (ES) cells, embryonal carcinoma (EC) cells, and embryonic germ (EG) cells. The expression of TRA-1-60 on human ES cells is down-regulated upon differentiation. The TRA-1-60 antibody recognizes a neuraminidase-resistant carbohydrate epitope expressed on podocalyxin, a member of the CD34-related family of sialomucins. Podocalyxin is a transmembrane glycoprotein, which has been implicated in the development of aggressiveness in a variety of cancers, including breast and prostate cancer.
<b>Catalog Number</b>	09-0010
<b>Size</b>	100 µl
<b>Concentration</b>	0.5 mg/ml
<b>Clone</b>	TRA-1-60
<b>Isotype</b>	Mouse IgM
<b>Immunogen</b>	Human embryonal carcinoma cell line 2102Ep
<b>Reactivity</b>	Human
<b>Preparation</b>	This antibody was purified by affinity chromatography.
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, and 0.09% sodium azide
<b>Storage and Stability</b>	Store at 4°C protected from light. Do not freeze. Stable for 6 months from date of receipt when stored as directed.
<b>Quality Control</b>	Tested by immunocytochemistry ( <b>Figure 1</b> ) and flow cytometry ( <b>Figure 2</b> ) to ensure product quality.
<b>Recommended Usage</b>	The suggested use of this antibody is a 1:100 dilution for immunocytochemistry and 0.25 µg per 1 x 10 <sup>6</sup> viable cells in 100 µl for flow cytometry. For application specific protocols, please reference <i>Protocol: Immunocytochemistry</i> and <i>Protocol: Flow Cytometry</i> online at <a href="http://www.stemgent.com/support/protocols">www.stemgent.com/support/protocols</a> .

For research use only. Not for use in diagnostic procedures.

## Product Specification Sheet



**Figure 1.** Immunocytochemistry analysis of TRA-1-60 on H1 human ES cells. Cells were stained with TRA-1-60 Antibody (Purified) at a 1:100 dilution followed by a secondary Cy<sup>™</sup> 3 antibody (red). DAPI staining was performed to visualize nuclei (blue).



**Figure 2.** Flow cytometry analysis of TRA-1-60 on H1 human ES cells. Red histogram represents TRA-1-60 Antibody (Purified) and open histogram represents isotype control at the same concentration.

### References

1. Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., and Yamanaka, S. (2007) Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors. *Cell* 131: 861-872.
2. Yu, J., Vodyanik, M.A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J.L., Tian, S., Nie, J., Jonsdottir, G.A., Ruotti, V., Stewart, R., Slukvin, I.I., and Thomson, J.A. (2007) Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318: 1917-1920.
3. Schopperle, W.M., and DeWolf, W.C. (2007) The TRA-1-60 and TRA-1-81 human pluripotent stem cell markers are expressed on podocalyxin in embryonal carcinoma. *Stem Cells* 25: 723-730.
4. Chin, A.C., Fong, W.J., Goh, L.T., Philp, R., Oh, S.K., and Choo, A.B. (2007) Identification of proteins from feeder conditioned medium that support human embryonic stem cells. *J Biotechnol.* 130: 320-328.
5. Badcock, G., Pigott, C., Goepel, J., and Andrews, P.W. (1999) The human embryonal carcinoma marker antigen TRA-1-60 is a sialylated keratan sulfate proteoglycan. *Cancer Res.* 59: 4715-4719.
6. Andrews, P.W., Banting, G., Damjanov, I., Arnaud, D., and Avner, P. (1984) Three monoclonal antibodies defining distinct differentiation antigens associated with different high molecular weight polypeptides on the surface of human embryonal carcinoma cells. *Hybridoma* 3: 347.

For research use only. Not for use in diagnostic procedures.